

REQUIRED IMMUNIZATIONS FOR SCHOOL ATTENDANCE

Kg-12th Grade

2007-2008

<http://www.dshs.state.tx.us/immunize/school/default.shtm>

Diphtheria, Tetanus, & Pertussis
(DTaP/DTP/DT/Tdap/Td)

Five doses of a diphtheria-tetanus-pertussis containing vaccine, one of which must have been received on or after the 4th birthday; however, 4 doses meet the requirement if the 4th dose was given on or after the 4th birthday.

Ages 7 years or older

(pertussis vaccine is not required)

Tdap/Td/DTP/DTaP or any combination of these

Three doses, including one dose on or after the 4th birthday

Booster Dose

Tdap/Td

One dose is required ten years after last dose of
DTP/DTaP/DT

Polio

Four doses of polio vaccine one of which must have been received on or after the 4th birthday; however, 3 doses meet the requirement if the 3rd dose was given on or after the 4th birthday

Measles, Mumps, and Rubella

Two doses of a measles-containing vaccine with the first dose on or after the 1st birthday; second dose required upon entry into kindergarten

Hepatitis B

Three doses

Varicella

One dose on or after 1st birthday
(Two doses if vaccine given at 13 years of age or older)

REQUIRED IMMUNIZATIONS FOR
PRE-K
2007-2008

In addition to the immunizations listed above; the following are required:

Hepatitis A	Two doses after 1 st birthday
Pneumococcal Conjugate	Four doses with one dose on or after 12 months of age OR: Three doses if a child received the 1 st dose from 7 through 11 months of age, and at least one dose was administered on or after 12 months of age OR: Two doses if a child received the 1 st dose from 12 Months through 23 months of age OR: One dose if a child is unvaccinated or partially vaccinated and healthy OR: If a child is unvaccinated or partially vaccinated and at high risk, two doses are required.
Haemophilus influenzae type b	Four doses, at least one received on or after 1 st birthday

RECOMMENDED IMMUNIZATIONS

Varicella	Second dose unless student has had chicken pox illness
Hepatitis A	Two doses
Meningococcal	One dose ages 11-18 years of age
Human Papillomavirus (HPV)	Three doses ages 9-26 years of age
Influenza (FLU)	One dose yearly